
INTERNATIONAL JOURNAL OF CURRENT RESEARCH IN BIOLOGY AND MEDICINE

ISSN: 2455-944X

www.darshanpublishers.com

DOI:10.22192/ijcrbm

Volume 2, Issue 3 - 2017

Original Research ArticleDOI: <http://dx.doi.org/10.22192/ijcrbm.2017.02.03.004>

In-vitro Anticancer activity of *Boerhaavia diffusa* Linn.

Dr R.Venkatajothi

Former, Senior Lecturer, Department of Microbiology, Faculty of Medicine, AIMST University, Kedah Darul Aman, Malaysia.

*Corresponding author: drvjothi10@gmail.com

Abstract

Cervical cancer is the second most common malignancy and cause of cancer mortality after breast cancer in developing countries. The high risk Human papilloma virus (HPV) types 16 and 18 are responsible for about 70% of all cervical cancer cases. HPV type 16 is sexually transmitted and most commonly causes Cervical cancer is malignant neoplasm of the cervix or cervical area. Modern research in the anticancer drug expansion from plants has been focused on investigate the molecular mechanism by which an agent influences cytotoxicity and apoptosis in cancer cells. The aim of this study is to analysis the prevalence of HPV type16 among women and to evaluate phytochemical investigations and cytotoxic activity of *Boerhaavia diffusa* Linn. A total of 1164 abnormal cervical specimens were tested for HPV type 16. Among of them HPV type 16 was the most prevalent type detected 528 (45.36%). The non HPV 16 types were 636 (54.63 %). The age wise distribution of HPV type 16 screening showed that 45.26% of patients were between the age group of 21-35, followed by 35.79% of the patients with the age group below 20 years. The overall in other group the prevalence ratio is generally lower. *B.diffusa* is a major medically importance weed, which has been proposed for the treatment of liver disorders, cancer and all types of inflammations. The active ingredients of *B. diffusa* were identified by ethanolic extraction method and further subjected to preliminary Phytochemical screening and Cytotoxicity activity by MTT assay. Phytochemical testing showed the presence of Alkaloids, Flavonoids, Tannins, Saponins, Terpenes, Anthraquinones and Steriods. Human Papiloma virus type 16 infected SiHa cervical cancer cell line were used to screen the anticancer potential of the extract of *B. diffusa*. The inhibition percentage with regard to cytotoxicity was found to be 96.3 % at 1000 µg/ml. The results of this research showed that the antioxidant played an important role in protecting the human body against cancer. The plant *Boerhaavia diffusa* Linn showed that significant anticancer activities against SiHa cell line which can be used as an alternative medicine.

Keywords: Cervical cancer, HPV type 16, *Boerhaavia diffusa*, Cytotoxicity

Introduction

Cervical cancer is one of the most commonly occurring cancers of the female reproductive tract¹⁵. Cervical cancer is the second most common cancer among women worldwide, with an estimated 529,409 new cases and 274,883 deaths in 2008¹². Worldwide in 2012, there are 528,000 cases of cervical cancer were estimated to have occurred, with 266,000 deaths¹⁷. An estimated 12,900 new cervical cancers and 4,100 cervical cancer deaths will occur in the

United States in 2015¹⁶. In Eastern and Middle Africa, cervical cancer is the most common cancer in women.

HPV infection has been shown to be an essential cause of cervical cancer. HPV can infect either cutaneous or mucosal epidermis. The high risk HPV types 16 and 18 are responsible for about 70% of all cervical cancer cases, followed by other HPV types in worldwide⁸.

Cervical cancer shows various types of symptoms and one of the most common symptoms is abnormal vaginal bleeding, but in some cases there may be no obvious symptoms until the cancer is in its advanced stages.

Cervical cancer is associated with many risk factors including early sexual debut, having multiple sexual partners or having sex with someone who has multiple sexual partners, HIV positive, a family history of cervical cancer, smoking and poverty etc. Cervical cancer ranks as the first most frequent cancer among women between 15 and 44 years of age in Tanzania.

Cervical cancer is one of the most common type of cancer in women and the one with the highest mortality rate⁷ in Tanzania. It has been reported that the high rate of infection with HPV type 16 and 18 are the frequent presence of viral DNA in cervical cancer among the population of Tanzanian women⁶. A recent estimate showed that approximately 7500 new cases are diagnosed every year and that about 4000 dies from cervical cancer yearly.

Boerhaavia diffusa Linn. commonly known as 'Punarnava' is an abundant creeping weed of the family *Nyctaginaceae* and it is commonly known as **Mukkurattai** in Tamil language, Hog Weed and Pig Weed in English. *B. diffusa* is a traditional herb of which has been proposed for the treatment of cancer, tumors, jaundice and liver disorders. The root is generally used as an infusion to treat internal inflammation. The whole plant extract is hepatoprotective in nature². It is also used for the treatment of diabetes¹⁰ and to treat seminal weakness and high blood pressure³.

The plant *B. diffusa* containing phenolic compounds, in particular alkaloids and amino acids have been reported to exhibit strong antioxidant properties. It also contains quinolizidine alkaloids and potassium salts¹³. The present investigation has been carried out to evaluate the cytotoxic effects of *B. diffusa* whole plant extracts. The whole plants of *Boerhaavia diffusa* are reported to have good medicinal values in traditional system of medicines.

Objectives of the Study

The aim of the present study is to analysis the prevalence of HPV type16 among women and identifies the inhibition of HPV 16 infected cervical cancer SiHa cell line by plant extract of *Boerhaavia diffusa* Linn.

Materials and Methods

A total of 1164 abnormal Pap smear specimens were tested for HPV type 16. The cases were women aged 18 to 65 years. Cervical cancer specimens obtained from women attending Cancer hospital in Tanzania, East Africa. They were recruited from the out and in-patient departments. The cases were women aged 18 years or more, with newly diagnosed histological confirmed cancer of the cervix.

Data will be analyzed by using SPSS. Cervical specimens were taken for cytological analysis. Cervical cancer was confirmed on the basis of histological results of cervical specimens and hybrid Capture 2 technology HPV DNA test. Human Papiloma virus type 16 infected SiHa cervical cancer cell line was obtained from National Centre for Cell Sciences (NCCS) at Pune in India.

The whole plant *B. diffusa* was collected, identified and authenticated by Dr. V. Balasubramanian, Senior Agriculture Officer at Salem in Tamil Nadu. The plant material was cut into small pieces and washed with sterile water, dried in shade, finely powdered and stored in air tight bottles. The active ingredients of *B. diffusa* were identified by ethanolic extraction method and further subjected to preliminary Phytochemical screening and Cytotoxicity activity by MTT assay. Phytochemical screening of plants was carried out to detect bioactive compounds using qualitative tests. Phytochemical test and MTT assay test done at Life Teck Research centre in Vadapalani at Chennai in India.

Inclusion criteria:

Inclusion criteria were based on the abnormal Pap smear test.

Exclusion criteria:

Exclusion criteria were normal Pap smear test.

Results and Discussion

A total of 1164 abnormal cervical specimens were tested for HPV type 16 (**Table1**). Among of them HPV type 16 was the most prevalent type detected 528 (45.36%). The non HPV 16 types were 636 (54.63 %). The age wise distribution of HPV type 16 screening showed that 45.26% of patients were between the age group of 21-35, followed by 35.79% of the patients with the age group below 20 years. 11.17% of them

belong to the age group of 36-40, 5.3% in the age group between 41- 49 and 2.46% between the age group of 50-60 (Table 2).

Table 1. Prevalence of HPV type 16 in women cases

S. No	Contents	Total numbers	Percentage (%)
1.	Total case study	1164	100%
2.	HPV type 16 Positive	528	45.36%
3.	Other than HPV type 16	636	54.63%

Table 2. Age wise distribution of HPV type 16 in Tanzanian women

S. No	Age group	HPV 16 Positive sample Size = 528	Percentage (%)
1.	Below 20	189	35.79%
2.	21 - 35	239	45.26%
3.	36 - 40	59	11.17%
4.	41 - 49	28	05.30%
5.	50 - 60	13	02.46%
6.	61 and above	0	0%

Among the total abnormal cervical specimens, 45% had HPV type 16 positive. Hence, the incidences of HPV type 16 were higher among Tanzanian women than the other types. Similarly, higher prevalence of HPV 16 were observed in Europe, with a few studies in Spain and Italy reporting type-specific prevalence of over 25% for HPV 16^{1,4,14}. This may be true in the present study also.

In the present findings clinically apparent genital HPV type 16 infections are correlated with age and suggesting that these viruses are primarily transmitted in adulthood through sexual intercourse⁵. It has been expected that at least 50% of sexually active men and women acquire genital HPV infection at some stage in their lives. Nearly 80% of women will have acquired

genital HPV by age 40 years, which makes HPV infections the norm rather than the exception suggested by Myers *et al*⁹. In Tanzania, screening is still at best very scarce and in the majority places not available at all.

The Phytochemical testing result showed that, the plant *B. diffusa* has contains major photochemicals such as Tannins, Alkaloids, Saponins, Flavonoids, Terpenes, Anthraquinones and Sterioids (Table 3) are well known natural antioxidant. Antioxidants thus play an important role in protecting the human body against damage by reactive oxygen species. Currently available drugs and synthetic drugs do have potential adverse reactions. There are many natural drugs which are yet to be explored systematically.

Table 3. Preliminary phytochemical analysis of Ethanol extract of *B. diffusa*

S. No	Test	Ethanol extract of <i>B. diffusa</i>
1.	Tannins Test -1 Test -2	Positive Positive
2.	Alkaloids	Positive
3.	Saponins	Positive
4.	Flavonoids	Positive
5.	Terpenes	Positive
6.	Anthroquinones	Positive
7.	Sterioids	Positive

The inhibition percentage with regard to cytotoxicity was found to be 96.3 % at 1000 µg/ml (Table 4). Rakhi Srivastava results demonstrated that *B. diffusa* fraction inhibits the proliferation of human cervical cancer cell line, HeLa and the cell cycle via S-phase

inhibition plays some roles in *B. diffusa* induced antiproliferative activities in the HeLa cell line¹¹. In the present study also showed that *B. diffusa* fraction inhibits SiHa cervical cancer cell line.


Table 4. Anticancer effect of *B. diffusa* on SiHa cell line

S. No	Concentration (µg/ml)	Dilutions	Absorbance (O.D)	Cell viability (%)
1.	1000	Neat	0.02	3.7
2.	500	1:1	0.11	20.7
3.	250	1:2	0.19	35.8
4.	125	1:4	0.25	47.1
5.	62.5	1:8	0.33	62.2
6.	31.2	1:16	0.40	75.4
7.	15.6	1:32	0.46	86.7
8.	7.8	1:64	0.50	94.3
9.	Cell control	-	0.53	100

Hence the plant *B. diffusa* has significant anticancer activities against SiHa cell line which can be used as an alternative medicine. The results of the study revealed that the plant extracts have strong antioxidant activity and remarkable cytotoxic activity (Fig. 1). The cytotoxic activity may be due to the presence of alkaloids in the plant.

Further investigations are needed to provide some additional insight into the in-vivo antioxidant activity and cytotoxic activity of the plant. The above results suggest that *Boerhaavia diffusa* Linn were found to reveal antioxidant potential which supports the use of this plant in cervical cancer treatment.

Figure 1. Anticancer effect of *B. diffusa* Linn on SiHA cell line (IC₅₀ Value)



Conclusion

This present study showed prevalence of HPV type 16 in women, is similar to those reported in other regions of the world. Similarly, it appears that HPV type 16 is the most common type associated with cervical

cancer. The above study shows those young groups are more vulnerable to HPV 16 virus infection. This will also require the treatment of male partner if infected. Currently less effective vaccinations such as Gardasil and Cervarix are available. The vaccines are useful only for women before exposure to HPV infection.

So, it is important to develop novel potent anti-cancer agents from natural products for the treatment of this virus disease. The plant *Boerhaavia diffusa* Linn has inhibition percentage with regard to cytotoxicity was found to be 96.3 % at 1000 µg/ml. The cytotoxic activity can be due to the presence of alkaloids in the plant, which needs further analysis. Further in-vivo studies will help us to investigation this plant in order to identify, isolate its active anticancer principle(s) and its mechanism of action.

References

1. Centurioni, M.G., Puppo, A. and Merlo, D.F. (2005). Prevalence of human papillomavirus cervical infection in an Italian asymptomatic population. *BMC Infect Dis*, 5: 77.
2. Evans WC. Trease and Evans Pharmacognosy. 14th Edn. W.B. Saunders: An imprint of Elsevier, London, 2002,437.
3. Gaitonde BB, Kulkarni HJ, Nabar SD. Diuretic activity of punarnava (*Boerhaavia diffusa*). 2nd ed. Bombay, India, 1974, 24.
4. Herbsleb, M., Knudsen, U.B. and Orntoft, T.F. (2001). Telomerase activity, MIB-1, PCNA, HPV 16 and p53 as diagnostic markers for cervical intraepithelial neoplasia. *APMIS*, 109(9): 607–17.
5. Ley, C., Bauer, H.M., Reingold, A., Schiffman, M.H., Chambers, J.C., Tashiro, C.J. and Manos, M.M. (1991). Determinants of genital human papillomavirus infection in young women. *Journal of the National Cancer Institute*, 83: 997-1003.
6. Mayaud P, Gill DK, Weiss HA, *et al.* The interrelation of HIV, cervical human papillomavirus, and neoplasia among antenatal clinic attenders in Tanzania. *Sex Transm Infect.* 2001;77:248-254.
7. Moscicki AB, Schiffman M, Kjaer S, Villa LL. Updating the natural history of HPV and anogenital cancer. *Vaccine*. 2006; 24: S3: 42-51.
8. Munoz, N., Bosch, F.X., de Sanjose, S., Herrero, R., Castellsague, X. and Shah, K.V. (2003). Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N Eng J Med*, 348: 518–27.
9. Myers ER, McCrory DC, Nanda K, Bastian L, Matchar DB. Mathematical model for the natural

- history of human papillomavirus infection and cervical carcinogenesis. *Am J Epidemiol* 2000; 151:1158-71.
10. Pari L, Amarnath SM. Antidiabetic activity of *Boerhaavia diffusa* L. effect on hepatic key enzymes in experimental diabetes. *J. Ethnopharmacol*, 2004, 91(1): 109-113.
 11. Rakhi Srivastava, Daman Saluja, Bilikere S. Dwarakanath, and Madhu Chopra. (2011). Inhibition of Human Cervical Cancer Cell Growth by Ethanolic Extract of *Boerhaavia diffusa* Linn. (Punarnava) Root. *Evidence-Based Complementary and Alternative Medicine*, 1-13.
 12. Sankaranarayanan R, Ferlay J. Worldwide burden of gynecological cancer: The size of the problem. *Best Pract Res Clin Obstet Gynaecol* 2006;20:207-25.
 13. Satheesh MA, Pari L. Antioxidant effect of *Boerhaavia diffusa* L. in tissues of alloxan induced diabetic rats. *Indian J Exp Biol*, 2004, 42(10): 989-992.
 14. Speich, N., Schmitt, C. and Bollmann, R. (2004). Human papillomavirus (HPV) study of 2916 cytological samples by PCR and DNA sequencing: genotype spectrum of patients from the west German area. *J Med Microbiol*, 53: 125–8.
 15. Stanley M: Pathology and epidemiology of HPV infection in females. *Gynecol Oncol* 2010; 117:S5.
 16. U.S. Cancer Statistics Working Group. United States Cancer Statistics: 1999–2013 Incidence and Mortality Web-based Report. Atlanta (GA): Department of Health and Human Services, Centers for Disease Control and Prevention, and National Cancer Institute; 2016.
 17. World Cancer Report 2014. World Health Organization. 2014. pp. Chapter 5.12.

Access this Article in Online



Website:

www.darshanpublishers.com

Subject:

[Medicinal Plants](#)

Quick Response
Code

How to cite this article:

R. Venkatajothi. (2017). *In-vitro* Anticancer activity of *Boerhaavia diffusa* Linn. *Int. J. Curr. Res. Biol. Med.* 2(3): 20-24.

DOI: <http://dx.doi.org/10.22192/ijcrbm.2017.02.03.004>